



Mathematics

Paper 1



(Non-Calculator)

Higher Tier

Mark Scheme

Edexcel GCSE

SET 3

Question	Working	Answer	Notes
Q1		$2^4 \times 13$	M1 Attempt at prime factor tree or other valid method A1 $2^4 \times 13$ oe
Q2	$\frac{9}{4} \times \frac{5}{3} = \frac{45}{12} = \frac{15}{4} = 3\frac{3}{4}$	$3\frac{3}{4}$	M1 $\frac{9}{4}$ and $\frac{5}{3}$ M1 Attempts to multiply their numerators and denominators A1 cao
Q3			B1 Perpendicular bisector of AB constructed (construction lines must be seen) B1 Circle or part circle, radius $4cm$ around B B1 Correct area shaded

Question	Working	Answer	Notes
Q4	$300 \div 3 = 100$ $100 \times 50p = £50$ $50 - 40 = 10$, profit = £10 $\frac{10}{40} \times 100 = 25\%$	25%	M1 $300 \div 3 = 100$ M1 $100 \times 50p = £50$ M1 $50 - 40 = 10$, profit = £10 A1 cao
Q5	$3.65 \times 10^5 = 365000$ $36.5 \times 10^{-2} = 0.365$	0.0365 36.5×10^{-2} 365 3.65×10^5	M1 365000 or 0.365 seen A1 cao
Q6	Angle $BHG = 90^\circ$ Interior angle of a hexagon: $\frac{4 \times 180}{6} = 120^\circ$ Angle $ABH = 120 - 90 = 30$ $90 = 3 \times 30$ so angle $BHG = 3 \times$ angle ABH		B1 Angle $BHG = 90^\circ$ M1 Attempt at a correct method to find the interior angle of a hexagon (= 120°) M1 <i>ft</i> Angle $ABH =$ their ' 120 ' $- 90$ B1 Concluding statement
Q7a	$\frac{1}{2} \times 4 \times 7 = 14$ $14 \times 10 = 140m^3$	$140m^3$	M1 Area of cross-section = $14m^2$ A1 cao
Q7b	$60 = \frac{F}{70}$ $F = 60 \times 70 = 4200N$ $P = \frac{4200}{40} = 105 N/m^2$	$105 N/m^2$	M1 $60 = \frac{F}{70}$ oe M1 $F = 4200N$ A1 cao

Question	Working	Answer	Notes																																			
Q8	Total height of boys: $6 \times 130 = 780$ Total height of girls: $4 \times 120 = 480$ Total height of 10 players: $780 + 480 = 1260$ Mean height of 10 players: $\frac{1260}{10} = 126$	Yes	M1 $6 \times 130 = 780$ and $4 \times 120 = 480$ M1 Adds their total heights to find the total of all players, 1260, or $1260 \div 10$ seen. B1 Correct conclusion with correct working																																			
Q9	<table border="1"><thead><tr><th>Sequence</th><th>1st term</th><th>2nd term</th><th>3rd term</th><th>4th term</th><th>5th term</th><th>nth term</th></tr></thead><tbody><tr><td>A</td><td>3</td><td>7</td><td>11</td><td>15</td><td>19</td><td>$4n - 1$</td></tr><tr><td>B</td><td>1</td><td>4</td><td>9</td><td>16</td><td>25</td><td>n^2</td></tr><tr><td>C</td><td>$\frac{1}{3}$</td><td>1</td><td>3</td><td>9</td><td>27</td><td>3^{n-2}</td></tr><tr><td>D</td><td>$\frac{2}{3}$</td><td>3</td><td>6</td><td>7</td><td>-2</td><td>$n^2 - 3^{n-2}$</td></tr></tbody></table>	Sequence	1st term	2nd term	3rd term	4th term	5th term	n th term	A	3	7	11	15	19	$4n - 1$	B	1	4	9	16	25	n^2	C	$\frac{1}{3}$	1	3	9	27	3^{n-2}	D	$\frac{2}{3}$	3	6	7	-2	$n^2 - 3^{n-2}$	19, 25 $4n - 1$ n^2 3^{n-2} $n^2 - 3^{n-2}$	A1 1st row correct A1 2nd row correct A1 3rd row correct A1 4th row correct
Sequence	1st term	2nd term	3rd term	4th term	5th term	n th term																																
A	3	7	11	15	19	$4n - 1$																																
B	1	4	9	16	25	n^2																																
C	$\frac{1}{3}$	1	3	9	27	3^{n-2}																																
D	$\frac{2}{3}$	3	6	7	-2	$n^2 - 3^{n-2}$																																
Q10a	$3 \times 4 = 12$ $12p^{2+3}q^{1+2} = 12p^5q^3$	$12p^5q^3$	M1 12 or p^5 or q^3 seen A1 cao																																			
Q10b	$a = 4(2m^2)^3$ $a = 4 \times 8m^6$ $a = 32m^6$	$a = 32m^6$	M1 $a = 4(2m^2)^3$ A1 cao																																			
Q10c	$32^{\frac{1}{2}} \times 2^{-2} = (2^5)^{\frac{1}{2}} \times 2^{-2}$ $= 2^{\frac{5}{2}} \times 2^{-2}$ $= 2^{\frac{1}{2}}$ Or $4\sqrt{2} \times \frac{1}{4} = \sqrt{2} = 2^{\frac{1}{2}}$	$2^{\frac{1}{2}}$	M1 Writes 32 as 2^5 or $4\sqrt{2}$ M1 Writes $\frac{1}{2^2}$ in a form which allows them to simplify the expression A1 cao written in index form																																			

Question	Working	Answer	Notes														
<div>Q11a</div>	<table><tr><th>Age (A years)</th><th>Cumulative frequency</th></tr><tr><td>$0 < A \leq 40$</td><td>13</td></tr><tr><td>$0 < A \leq 80$</td><td>41</td></tr><tr><td>$0 < A \leq 120$</td><td>64</td></tr><tr><td>$0 < A \leq 160$</td><td>73</td></tr><tr><td>$0 < A \leq 200$</td><td>78</td></tr><tr><td>$0 < A \leq 240$</td><td>80</td></tr></table>	Age (A years)	Cumulative frequency	$0 < A \leq 40$	13	$0 < A \leq 80$	41	$0 < A \leq 120$	64	$0 < A \leq 160$	73	$0 < A \leq 200$	78	$0 < A \leq 240$	80		A1 all cumulative frequencies correct
Age (A years)	Cumulative frequency																
$0 < A \leq 40$	13																
$0 < A \leq 80$	41																
$0 < A \leq 120$	64																
$0 < A \leq 160$	73																
$0 < A \leq 200$	78																
$0 < A \leq 240$	80																
<div>Q11b</div>			M1 At least 4 points plotted correctly A1 All points correct and joined with an appropriate curve														

Question	Working	Answer	Notes
<p>Q11c</p>	<p>54 trees less than 100 years old $80 - 54 = 26$ trees over 100 years old</p>	$\frac{26}{80}$	<p>M1 Line drawn at 100 years and value of 25 - 27, read from their graph</p> <p>M1 Subtracts their '26' from 80</p> <p>A1 $\frac{26}{80}$ oe</p>
<p>Q12</p>	$\frac{28}{36} : \frac{22}{40} : \frac{16}{30} = \frac{280}{360} : \frac{198}{360} : \frac{192}{360}$ $= 280 : 198 : 192$ $= 140 : 99 : 96$	$140 : 99 : 96$	<p>M1 $\frac{28}{36} \frac{22}{40}, \frac{16}{30}$</p> <p>M1 Writes fractions with common denominator</p> <p>A1 cao</p>

Question	Working	Answer	Notes
Q13	$x = 0.4353535\dots$ $10x = 4.353535\dots$ $1000x = 435.353535\dots$ $1000x - 10x = 431$ $990x = 431$ $x = \frac{431}{990}$	$\frac{431}{990}$	M1 $10x$ and $1000x$ or $4.\dot{3}\ddot{5}$ and $435.\dot{3}\ddot{5}$ seen M1 Attempt to subtract $10x$ from $1000x$ or $4.\dot{3}\ddot{5}$ from $435.\dot{3}\ddot{5}$ A1 Correct fraction with correct steps shown
Q14	$\sin(30) = \frac{1}{2}, \cos(30) = \frac{\sqrt{3}}{2}$ $6 \times \frac{1}{2} \times 2 \times \frac{\sqrt{3}}{2} = 6 \times \frac{\sqrt{3}}{2} = 3\sqrt{3}$	$a = 3$	M1 $\sin(30) = \frac{1}{2}, \cos(30) = \frac{\sqrt{3}}{2}$ seen or implied M1 $6 \times \frac{1}{2} \times 2 \times \frac{\sqrt{3}}{2}$ A1 cao
Q15	$0.6 \times 0.4 + 0.4 \times 0.1 = 0.24 + 0.04$ $= 0.28$	0.28	M1 Tree diagram drawn or attempt at ‘walks and late’ and ‘bus and late’ M1 $0.6 \times 0.4 = 0.24$ M1 $0.4 \times 0.1 = 0.04$ A1 cao
Q16	$P = \frac{k}{Q^2}$ $1.5 = \frac{k}{10^2}, k = 1.5 \times 100 = 150$ $P = \frac{150}{5^2} = \frac{150}{25} = 6$	$P = 6$	M1 $1.5 = \frac{k}{10^2}$ M1 $k = 150$ A1 cao

Question	Working	Answer	Notes
Q17	$(x + 5)^2 + (x - 2)^2 = (x + 6)^2$ $x^2 + 10x + 25 + x^2 - 4x + 4 = x^2 + 12x + 36$ $x^2 - 6x - 7 = 0$ $(x + 1)(x - 7) = 0$ $x = -1$ (invalid) or $x = 7$	$x = 7$	M1 $(x + 5)^2 + (x - 2)^2 = (x + 6)^2$ oe M1 $x^2 - 6x - 7 = 0$ M1 Valid attempt to solve their quadratic A1 cao (must disregard $x = -1$)
Q18	$b = -6$ $0 = 1^2 + a - 6, a = 5$ $y = x^2 + 5x - 6$ $y = (x + \frac{5}{2})^2 - (\frac{5}{2})^2 - 6$ $y = (x + \frac{5}{2})^2 - \frac{49}{4}$	$(-\frac{5}{2}, -\frac{49}{4})$	M1 $b = -6$ M1 $0 = 1^2 + a - 6, a = 5$ or attempt to identify second factor bracket e.g. $(x - 1)(x + 6)$ M1 Completes square or uses other method to determine x coordinate $= -\frac{5}{2}$ or -2.5 A1 oe e.g. $(-2.5, -12.25)$
Q19	$x = 3y$ $2x + 2z = 26 \Rightarrow 6y + 2z = 26$ $y + 2z = 11$ $5y = 15$ $y = 3$ $x = 3 \times 3 = 9$ $3 + 2z = 11$ $z = 4$	$x = 9$ $y = 3$ $z = 4$	M1 $2x + 2z = 26$ and $y + 2z = 11$ M1 $x = 3y$ and substitutes into one of the first to equations to get 2 equations in 2 variables M1 Valid attempt to solve simultaneous equations M1 $x = 9$ or $y = 3$ or $z = 4$ seen A1 cao

Question	Working	Answer	Notes
Q20	$M = \frac{\sqrt{90}}{2 + \sqrt{10}} = \frac{3\sqrt{10}}{2 + \sqrt{10}}$ $= \frac{3\sqrt{10}(2 - \sqrt{10})}{(2 + \sqrt{10})(2 - \sqrt{10})}$ $= \frac{6\sqrt{10} - 30}{4 - 10}$ $= \frac{6\sqrt{10} - 30}{-6}$ $= 5 - \sqrt{10}$	$5 - \sqrt{10}$	<p>M1 $M = \frac{\sqrt{90}}{2 + \sqrt{10}}$</p> <p>M1 attempt to rationalise the denominator, e.g., multiplying by $\frac{2 - \sqrt{10}}{2 - \sqrt{10}}$</p> <p>M1 Reaches $\frac{6\sqrt{10} - 30}{-6}$</p> <p>A1 cao</p>
Q21a	$A = h \times b \Rightarrow 54 = h \times b$ $h = \frac{54}{b}$		A1 cao
Q21b	$2h + 2b = 33$ $2\left(\frac{54}{b}\right) + 2b = 33$ $108 + 2b^2 = 33b$ $2b^2 - 33b + 108 = 0$ $(2b - 9)(b - 12) = 0$ $b = \frac{9}{2} \text{ or } b = 12$	$\frac{9}{2}$ and 12	<p>M1 $2h + 2b = 33$ and substitutes $\frac{54}{b}$</p> <p>or $2b^2 - 33b + 108 = 0$ seen</p> <p>M1 Valid attempt to solve quadratic</p> <p>A1 cao</p>

Help ease the pressure with a personalised revision programme for each of your target KS4 students

Our one to one GCSE revision programme is designed to help your target students reach their potential in their GCSE maths exams.

Our specialist maths tutors work one to one with each student, focusing on securing core KS4 content and building familiarity with the kinds of questions they'll be tackling in their GCSE exams.

Get in touch today:

✉ hello@thirdspacelearning.com

🔍 thirdspacelearning.com

☎ 0203 771 0095