



# Mathematics

## Paper 5


### (Non-Calculator)

### Higher Tier

### Mark Scheme

OCR GCSE

SET 3

Question	Working	Answer	Notes
Q1	$\frac{9}{4} \times \frac{5}{3} = \frac{45}{12} = \frac{15}{4} = 3\frac{3}{4}$	$3\frac{3}{4}$	M1 $\frac{9}{4}$ and $\frac{5}{3}$ M1 Attempts to multiply their numerators and denominators A1 cao
Q2a	<div><p><math>2^4 \times 13</math></p></div>	4	A1 cao
Q2b	$78 = 2 \times 3 \times 13$ $HCF: 2 \times 13 = 26$	26	M1 78 written as product of primes OR at least 6 correct factors of 78 listed M1 Identifies common prime factors OR identifies a common factor A1 cao

Question	Working	Answer	Notes
Q3			B1 Perpendicular bisector of $AB$ constructed (construction lines must be seen) B1 Circle or part circle, radius $4cm$ around $B$ B1 Correct area shaded
Q4	$2 \text{ litres} = 2000ml$ $75\% \text{ of } 2000ml: 2000 \div 4 = 500,$ $500 \times 3 = 1500ml$ $240 \times 6 = 1440ml$	No	M1 $240 \times 6 = 1440ml$ M1 $2 \text{ litres} = 2000ml$ seen or implied M1 $75\% \text{ of } 2000 = 1500ml$ OR $\frac{1440}{2000} \times 100 = \frac{72}{100} \times 100 = 72\%$ A1 Correct answer following correct working
Q5	$300 \div 3 = 100$ $100 \times 50p = \text{£}50$ $50 - 40 = 10, \text{ profit} = \text{£}10$ $\frac{10}{40} \times 100 = 25\%$	25%	M1 $300 \div 3 = 100$ or $100 \times 50p = \text{£}50$ M1 $50 - 40 = 10, \text{ profit} = \text{£}10$ M1 Attempt to calculate the percentage profit A1 cao
Q6	$3.65 \times 10^5 = 365000$ $36.5 \times 10^{-2} = 0.365$	0.0365 $36.5 \times 10^{-2}$ 365 $3.65 \times 10^5$	M1 365000 or 0.365 seen A1 cao

Question	Working	Answer	Notes
<b>Q7</b>	Angle $BHG = 90^\circ$ Interior angle of a hexagon: $\frac{4 \times 180}{6} = 120^\circ$ Angle $ABH = 120 - 90 = 30$ $90 = 3 \times 30$ so angle $BHG = 3 \times$ angle $ABH$		B1 Angle $BHG = 90^\circ$ M1 Attempt at a correct method to find the interior angle of a hexagon M1 <i>ft</i> Angle $ABH =$ their '120' $- 90$ B1 Concluding statement
<b>Q8a</b>	Volume estimate: $10 \times 10 \times 10 = 1000$ Density estimate: $\frac{600}{1000} = 0.6g/cm^3$	$0.6g/cm^3$	M1 Rounds 10.1 to 10 M1 Rounds 594 to 600 M1 $10 \times 10 \times 10 = 1000$ A1 cao
<b>Q8b</b>		Over-estimate	B1
<b>Q9a</b>	$\frac{1}{2} \times 4 \times 7 = 14$ $14 \times 10 = 140m^3$	$140m^3$	M1 Area of cross-section = $14m^2$ A1 cao
<b>Q9b</b>	$60 = \frac{F}{70}$ $F = 60 \times 70 = 4200N$ $P = \frac{4200}{40} = 105 \text{ newtons}/m^2$	$105 \text{ newtons}/m^2$	M1 $60 = \frac{F}{70}$ oe M1 $F = 4200N$ A1 cao



Question	Working	Answer	Notes
<b>Q10</b>	Total height of boys: $6 \times 130 = 780$ Total height of girls: $4 \times 120 = 480$ Total height of 10 players: $780 + 480 = 1260$ Mean height of 10 players: $\frac{1260}{10} = 126$	Yes	M1 $6 \times 130 = 780$ or $4 \times 120 = 480$ M1 $780 + 480 (= 1260)$ M1 $1260 \div 10$ A1 Concludes 'Yes' with correct working
<b>Q11a</b>	$3 \times 4 = 12$ $12p^{2+3} q^{1+2} = 12p^5 q^3$	$12p^5 q^3$	M1 12 or $p^5$ or $q^3$ seen A1 cao
<b>Q11b</b>	$a = 4(2m^2)^3$ $a = 4 \times 8m^6$ $a = 32m^6$	$a = 32m^6$	M1 $a = 4(2m^2)^3$ A1 cao
<b>Q11c</b>	$32^{\frac{1}{2}} \times 2^{-2} = (2^5)^{\frac{1}{2}} \times 2^{-2}$ $= 2^{\frac{5}{2}} \times 2^{-2}$ $= 2^{\frac{1}{2}}$ Or $4\sqrt{2} \times \frac{1}{4} = \sqrt{2} = 2^{\frac{1}{2}}$	$2^{\frac{1}{2}}$	M1 Writes 32 as $2^5$ or $4\sqrt{2}$ M1 Writes $\frac{1}{2^2}$ in a form which allows them to simplify the expression A1 cao written in index form

Question	Working	Answer	Notes														
Q12a	<table><thead><tr><th>Age (<math>A</math> years)</th><th>Cumulative frequency</th></tr></thead><tbody><tr><td><math>0 &lt; A \leq 40</math></td><td>13</td></tr><tr><td><math>0 &lt; A \leq 80</math></td><td>41</td></tr><tr><td><math>0 &lt; A \leq 120</math></td><td>64</td></tr><tr><td><math>0 &lt; A \leq 160</math></td><td>73</td></tr><tr><td><math>0 &lt; A \leq 200</math></td><td>78</td></tr><tr><td><math>0 &lt; A \leq 240</math></td><td>80</td></tr></tbody></table>	Age ( $A$ years)	Cumulative frequency	$0 < A \leq 40$	13	$0 < A \leq 80$	41	$0 < A \leq 120$	64	$0 < A \leq 160$	73	$0 < A \leq 200$	78	$0 < A \leq 240$	80		A1
Age ( $A$ years)	Cumulative frequency																
$0 < A \leq 40$	13																
$0 < A \leq 80$	41																
$0 < A \leq 120$	64																
$0 < A \leq 160$	73																
$0 < A \leq 200$	78																
$0 < A \leq 240$	80																
Q12b			M1 At least 4 points plotted correctly A1 All points correct and joined with an appropriate curve														

Question	Working	Answer	Notes
<p>Q12c</p>	<p>54 trees less than 100 years old <math>80 - 54 = 26</math> trees over 100 years old</p>	$\frac{26}{80}$	<p>M1 Line drawn at 100 years and value of 53 - 55, read from their graph</p> <p>M1 Subtracts their '54' from 80</p> <p>A1 <math>\frac{26}{80}</math> oe</p>
<p>Q13</p>	$\frac{28}{36} : \frac{22}{40} : \frac{16}{30} = \frac{280}{360} : \frac{198}{360} : \frac{192}{360}$ $= 280 : 198 : 192$ $= 140 : 99 : 96$	$140 : 99 : 96$	<p>M1 <math>\frac{28}{36}, \frac{22}{40}, \frac{16}{30}</math></p> <p>M1 Writes fractions with common denominator</p> <p>A1 cao</p>

Question	Working	Answer	Notes
Q14			M1 Line $y = x$ drawn with solid line M1 Line $y = x$ drawn with dotted line B1 A region which correctly satisfies at least one of the inequalities identified B1 A region which correctly satisfies at least two of the inequalities identified A1 Fully correct solution
Q15a	$5 \div 6 = 0.83333....$	$0.83333$ or $0.8\dot{3}$	M1 Attempt to divide 5 by 6 A1 cao
Q15b	$x = 0.4353535...$ $10x = 4.353535...$ $1000x = 435.353535....$ $1000x - 10x = 431$ $990x = 431$ $x = \frac{431}{990}$	$\frac{431}{990}$	M1 $10x$ and $1000x$ seen M1 Attempt to subtract $10x$ from $1000x$ A1 Correct fraction with correct steps shown

Question	Working	Answer	Notes
<b>Q16</b>	$\frac{\theta}{360} \times \pi \times 3^2 = 2$ $\frac{9\theta}{360} = 2$ $\theta = \frac{720}{9} = 80^\circ$ $\text{Arc length: } \frac{80}{360} \times 2 \times \pi \times 3 = \frac{4}{3} \pi$ $\text{Perimeter: } 6 + \frac{4}{3} \pi$	$6 + \frac{4}{3} \pi \text{ cm}$	<p>M1 <math>\frac{\theta}{360} \times \pi \times 3^2 = 2\pi</math></p> <p>M1 <math>\theta = \frac{720}{9} = 80^\circ</math></p> <p>M1 <i>ft</i> Arc length: <math>\frac{\text{their '80'}}{360} \times 2 \times \pi \times 3</math></p> <p>M1 Arc length = <math>\frac{4}{3} \pi</math></p> <p>M1 <i>ft</i> their '<math>\frac{4}{3} \pi</math>' + 6</p> <p>A1 cao</p>
<b>Q17</b>	$\sin(30) = \frac{1}{2}, \cos(30) = \frac{\sqrt{3}}{2}$ $6 \times \frac{1}{2} \times 2 \times \frac{\sqrt{3}}{2} = 6 \times \frac{\sqrt{3}}{2} = 3\sqrt{3}$	$a = 3$	<p>M1 <math>\sin(30) = \frac{1}{2}, \cos(30) = \frac{\sqrt{3}}{2}</math> seen or implied</p> <p>M1 <math>6 \times \frac{1}{2} \times 2 \times \frac{\sqrt{3}}{2} = 3\sqrt{3}</math></p> <p>A1 cao</p>
<b>Q18</b>	$0.6 \times 0.4 + 0.4 \times 0.1 = 0.24 + 0.04$ $= 0.28$	0.28	<p>M1 Tree diagram drawn or attempt at 'walks and late' and 'bus and late'</p> <p>M1 <math>0.6 \times 0.4 = 0.24</math></p> <p>M1 <math>0.4 \times 0.1 = 0.04</math></p> <p>A1 cao</p>
<b>Q19</b>	$P = \frac{k}{Q^2}$ $1.5 = \frac{k}{10^2}, k = 1.5 \times 100 = 150$ $P = \frac{150}{5^2} = \frac{150}{25} = 6$	$P = 6$	<p>M1 <math>1.5 = \frac{k}{10^2}</math></p> <p>M1 <math>k = 150</math></p> <p>A1 cao</p>

Question	Working	Answer	Notes
<b>Q20</b>	$(x + 5)^2 + (x - 2)^2 = (x + 6)^2$ $x^2 + 10x + 25 + x^2 - 4x + 4 = x^2 + 12x + 36$ $x^2 - 6x - 7 = 0$ $(x + 1)(x - 7) = 0$ $x = -1$ (invalid) or $x = 7$	$x = 7$	M1 $(x + 5)^2 + (x - 2)^2 = (x + 6)^2$ M1 $x^2 - 6x - 7 = 0$ M1 Valid attempt to solve their quadratic A1 cao (must disregard $x = -1$ )
<b>Q21</b>	$b = -6$ $0 = 1^2 + a - 6, a = 5$ $y = x^2 + 5x - 6$ $y = (x + \frac{5}{2})^2 - (\frac{5}{2})^2 - 6$ $y = (x + \frac{5}{2})^2 - \frac{49}{4}$	$(-\frac{5}{2}, -\frac{49}{4})$	M1 $b = -6$ M1 $0 = 1^2 + a - 6, a = 5$ M1 Correct method to determine $x$ coordinate $= -\frac{5}{2}$ A1 cao
<b>Q22</b>	$x = 3y$ $2x + 2z = 26 \Rightarrow 6y + 2z = 26$ $y + 2z = 11$ $5y = 15$ $y = 3$ $x = 3 \times 3 = 9$ $3 + 2z = 11$ $z = 4$	$x = 9$ $y = 3$ $z = 4$	M1 $2x + 2z = 26$ M1 $y + 2z = 11$ M1 $x = 3y$ and substitutes into one of the first to equations to get 2 equations in 2 variables M1 Valid attempt to solve simultaneous equations A1 cao

Question	Working	Answer	Notes
<b>Q23</b>	$M = \frac{\sqrt{90}}{2 + \sqrt{10}} = \frac{3\sqrt{10}}{2 + \sqrt{10}}$ $= \frac{3\sqrt{10}(2 - \sqrt{10})}{(2 + \sqrt{10})(2 - \sqrt{10})}$ $= \frac{6\sqrt{10} - 30}{4 - 10}$ $= \frac{6\sqrt{10} - 30}{-6}$ $= 5 - \sqrt{10}$	$5 - \sqrt{10}$	<p>M1 <math>M = \frac{\sqrt{90}}{2 + \sqrt{10}}</math></p> <p>M1 Multiplies by <math>\frac{2 - \sqrt{10}}{2 - \sqrt{10}}</math></p> <p>M1 Reaches <math>\frac{6\sqrt{10} - 30}{-6}</math></p> <p>A1 cao</p>
<b>Q24a</b>	$A = h \times b \Rightarrow 54 = h \times b$ $h = \frac{54}{b}$		A1 correct steps shown
<b>Q25b</b>	$2h + 2b = 33$ $2\left(\frac{54}{b}\right) + 2b = 33$ $108 + 2b^2 = 33b$ $2b^2 - 33b + 108 = 0$ $(2b - 9)(b - 12) = 0$ $b = \frac{9}{2} \text{ or } b = 12$	$\frac{9}{2}$ and 12	<p>M1 <math>2h + 2b = 33</math> and substitutes <math>\frac{54}{b}</math></p> <p>M1 Reaches <math>2b^2 - 33b + 108 = 0</math></p> <p>A1 cao</p>

# Help ease the pressure with a personalised revision programme for each of your target KS4 students

Our one to one GCSE revision programme is designed to help your target students reach their potential in their GCSE maths exams.

Our specialist maths tutors work one to one with each student, focusing on securing core KS4 content and building familiarity with the kinds of questions they'll be tackling in their GCSE exams.

Get in touch today:

✉ [hello@thirdspacelearning.com](mailto:hello@thirdspacelearning.com)

🔍 [thirdspacelearning.com](https://thirdspacelearning.com)

☎ 0203 771 0095